

A R C T I C O C E A N

MAP of
SEDIMENTARY BASINS of OFFSHORE ALASKA

FROM RESEARCH STUDIES BY

T.R. Bruns - Eastern Gulf of Alaska	M.A. Fisher - Lower Cook Inlet
R. von Huene - Western Gulf of Alaska	D.W. Schell - Bering Sea
T.R. Bruns - Chukchi, Beaufort Seas	M.S. Marlow - Hope Basin
M.A. Fisher - Norton Sound	A.K. Cooper
A. Grants	S.L. Eittreim
S.L. Eittreim	A. Grants

COMPILED BY
T.E. Chase, C.P. Miller, J.A. Thomas, B.A. Seekins, J.D. Young, and G.E. Carpenter

1979
SCALE
1:2,707,702 of Latitude 62°

KILOMETERS
NAUTICAL MILES

MERCATOR PROJECTION

NOTE:
THIS CHART IS A COMPILATION OF KNOWN SEDIMENTARY BASINS OF OFFSHORE ALASKA. IT IS INTENDED TO GIVE USERS A WORKING MAP FOR PLANNING PURPOSES. PLOTTING OF ADDITIONAL DATA, OR AID IN LOCATION OF SEDIMENTARY FEATURES.

LEGEND

Isopachs: Variable contour interval in kilometers
 Faults: Normal ——— Thrust ———
 Crest of arch ———
 Anticline ———
 Syncline ———
 Inferred boundary:

Base map from
NAVOCEAO No. 532
NAVOCEAO No. 520
NOS No. 16204

Bathymetric contours from

J. Greenberg	BEAUFORT SEA
P.E. Hart	
A. Grants	BERING SEA
D.W. Schell	
E.C. Buffington	NORTH PACIFIC OCEAN
D.M. Hopkins	
T.R. Alphe	
G.M. Schumacher	
T.E. Chase	
W.W. Menard	
J. Mommericks	

CANADA
SELECTED PROFILES

